

Bridge Simulation and Metric Estimation on Lie Groups

Mathias Højgaard Jensen¹, Sarang Joshi², and Stefan Sommer¹

¹ Department of Computer Science, University of Copenhagen
Universitetsparken 1, 2100, Copenhagen Ø, Denmark
{matje,sommer}@di.ku.dk

² Department of Biomedical Engineering, University of Utah
72 S Central Campus Drive, Salt Lake City, UT 84112, USA
sjoshi@sci.utah.edu

Abstract. We present a simulation scheme for simulating Brownian bridges on complete and connected Lie groups. We show how this simulation scheme leads to absolute continuity of the Brownian bridge measure with respect to the guided process measure. This result generalizes the Euclidean result of Delyon and Hu to Lie groups. We present numerical results of the guided process in the Lie group $SO(3)$. In particular, we apply importance sampling to estimate the metric on $SO(3)$ using an iterative maximum likelihood method.

Keywords: Brownian Motion, Brownian Bridge Simulation, Importance sampling, Lie groups, Metric estimation

1 Introduction

Bridge simulation techniques are known to play a fundamental role in statistical inference for diffusion processes. Diffusion bridges in manifolds have mainly been used to provide gradient and hessian estimates. To the best of our knowledge, this paper is the first to describe a simulation technique for diffusion bridges in the context of Lie groups.

The paper is organized as follows. In Section 2, we describe some background theory of Lie groups, Brownian motions, and Brownian bridges in Riemannian manifolds. Section 3 presents the theory and results. Section 4 shows in practice the simulation scheme in the Lie group $SO(3)$. Using importance sampling, we obtain an estimate of the underlying unknown metric.

2 Notation and Background

Lie Groups Throughout, we let G denote a connected Lie Group of dimension d , i.e., a smooth manifold with a group structure such that the group operations $G \times G \ni (x, y) \mapsto xy \in G$ and $G \ni x \mapsto x^{-1} \in G$ are smooth maps. If $x \in G$, the left-multiplication map, $L_x y$, defined by $y \mapsto \mu(x, y)$, is a diffeomorphism from G to itself. Similarly, the right-multiplication map $R_x y$ defines a diffeomorphism from G to itself by $y \mapsto \mu(y, x)$. We assume throughout that G acts on itself by left-multiplication. Let $dL_x: TG \rightarrow TG$ denote the pushforward map given by $(dL_x)_y: T_y G \rightarrow T_{xy} G$. A vector field V on G is said to be left-invariant if

$(dL_x)_y V(y) = V(xy)$. The space of left-invariant vector fields is linearly isomorphic to $T_e G$, the tangent space at the identity element $e \in G$. By equipping the tangent space $T_e G$ with the Lie bracket we can identify the Lie algebra \mathcal{G} with $T_e G$. The group structure of G makes it possible to define an action of G on its Lie algebra \mathcal{G} . The conjugation map $C_x := L_x \circ R_x^{-1} : y \mapsto xyx^{-1}$, for $x \in G$, fixes the identity e . Its pushforward map at e , $(dC_x)_e$, is then a linear automorphism of \mathcal{G} . Define $\text{Ad}(x) := (dC_x)_e$, then $\text{Ad} : x \mapsto \text{Ad}(x)$ is the adjoint representation of G in \mathcal{G} . The map $G \times \mathcal{G} \ni (x, v) \mapsto \text{Ad}(x)v \in \mathcal{G}$ is the adjoint action of G on \mathcal{G} . We denote by $\langle \cdot, \cdot \rangle$ a Riemannian metric on G . The metric is said to be left-invariant if $\langle u, v \rangle_y = \langle (dL_x)_y u, (dL_x)_y v \rangle_{L_x(y)}$, for every $u, v \in T_y G$, i.e., the left-multiplication maps are isometries, for every $x \in G$. In particular, we say that the metric is $\text{Ad}(G)$ -invariant if $\langle u, v \rangle_e = \langle \text{Ad}(x)u, \text{Ad}(x)v \rangle_e$, for every $u, v \in \mathcal{G}$. Note that an $\text{Ad}(G)$ -invariant inner on \mathcal{G} induces a bi-invariant (left- and right-invariant) metric on G .

Brownian Motion Endowing a smooth manifold M with a Riemannian metric, g , allows us to define the Laplace-Beltrami operator, $\Delta_M f = \text{div grad } f$. This operator is the generalization of the Euclidean Laplacian operator to manifolds. In terms of local coordinates (x_1, \dots, x_d) the expression for the Laplace-Beltrami operator becomes $\Delta_M f = \det(g)^{-1/2} \left(\frac{\partial}{\partial x_j} g^{ji} \det(g)^{1/2} \frac{\partial}{\partial x_i} \right) f$, where $\det(g)$ denotes the determinant of the Riemannian metric g and g^{ij} are the coefficients of the inverse of g . An application of the product rule implies that Δ_M can be rewritten as $\Delta_M f = a^{ij} \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} f + b^k \frac{\partial}{\partial x_j} f$, where $a^{ij} = g^{ij}$, $b^k = -g^{ij} \Gamma_{ij}^k$, and Γ denote the Christoffel symbols related to the Riemannian metric. This diffusion operator defines a Brownian motion on the M , valid up to its first exit time of the local coordinate chart.

In the case of the Lie group G , the identification of the space of left-invariant vector fields with the Lie algebra \mathcal{G} allows for a global description of Δ_G . Indeed, let $\{v_1, \dots, v_d\}$ be an orthonormal basis of $T_e G$. Then $V_i(x) = (dL_x)_e v_i$ defines left-invariant vector fields on G and the Laplace-Beltrami operator can be written as (cf. [6, Proposition 2.5]) $\Delta_G f(x) = \sum_{i=1}^d V_i^2 f(x) - V_0 f(x)$, where $V_0 = \sum_{i,j=1}^d C_{ij}^j V_j$ and C_{ij}^k denote the structure coefficients given by $[V_i, V_j] = C_{ij}^k V_k$. The corresponding stochastic differential equation (SDE) for the Brownian motion on G , in terms of left-invariant vector fields, then becomes

$$dX_t = -\frac{1}{2} V_0(X_t) dt + V_i(X_t) \circ dB_t^i, \quad X_0 = e, \quad (1)$$

where \circ denotes integration in the Stratonovich sense. By [6, Proposition 2.6], if the inner product is $\text{Ad}(G)$ invariant, then $V_0 = 0$. The solution of (1) is conservative or non-explosive and is called the left-Brownian motion on G (see [8] and references therein).

Riemannian Brownian Bridges In this section, we briefly review some classical facts on Brownian bridges on Riemannian manifolds. As Lie groups themselves are manifolds, the theory carries over *mutatis mutandis*. However, Lie groups' group structure allows the notion of left-invariant vector fields. The identification of the Lie algebra with the vector space of left-invariant vector fields

makes Lie groups parallelizable. Thus, the frame bundle construction for developing stochastic processes on manifolds becomes superfluous since left-invariant vector fields ensure stochastic parallel displacement.

Let \mathbb{P}_x^t be the measure of a Riemannian Brownian motion, X_t , at some time t started at point x . Suppose p denotes the transition density of the Riemannian Brownian motion. In that case, $d\mathbb{P}_x^t = p(t, x, y)d\text{Vol}(y)$ describes the measure of the Riemannian Brownian motion, where $d\text{Vol}(y)$ is the Riemannian volume measure. Conditioning the Riemannian Brownian motion to hit some point v at time $T > 0$ results in a Riemannian Brownian bridge. Here, $\mathbb{P}_{x,v}^T$ denotes the corresponding probability measure. The two measures are equivalent over the time interval $[0, T)$, however mutually singular at time $t = T$. The initial enlargement of the filtration remedies the singularity. The corresponding Radon-Nikodym derivative is given by

$$\frac{d\mathbb{P}_{x,v}^T}{d\mathbb{P}_x^T} \Big|_{\mathcal{F}_s} = \frac{p(T-s, X_s, v)}{p(T, x, v)} \quad \text{for } 0 \leq s < T,$$

which is a martingale for $s < T$. The Radon-Nikodym derivative defines the density for the change of measure and provides the basis for the description of Brownian bridges. In particular, it provides the conditional expectation defined by

$$\mathbb{E}[F(X_t)|X_T = v] = \frac{\mathbb{E}[p(T-t, X_t, v)F(X_t)]}{p(T, x, v)},$$

for any bounded and \mathcal{F}_s -measurable random variable $F(X_s)$. As described in [3], the Brownian bridge yields an SDE in the frame bundle, \mathcal{FM} , given by

$$dU_t = H_i(U_t) \circ (dB_t^i + H_i \log \tilde{p}(T-t, U_t, v)dt), \quad U_0 = u_0, \quad (2)$$

in terms of the horizontal vector fields (H_i), which is the lifted M -valued Brownian bridge, $X_t := \pi(U_t)$, where $\pi: \mathcal{FM} \rightarrow M$.

3 Simulation of Bridges on Lie Groups

In this section, we consider the task of simulating (1) conditioned to hit $v \in G$, at time $T > 0$. The potentially intractable transition density for the solution of (1) inhibits simulation directly from (2). Instead, we propose to add a guiding term mimicking that of Delyon and Hu [2], i.e., the guiding term becomes the gradient of the distance to v divided by the time to arrival. The SDE for the guided diffusion becomes

$$dY_t = -\frac{1}{2}V_0(Y_t)dt + V_i(Y_t) \circ \left(dB_t^i - \frac{(\nabla_y d(Y_t, v))^2}{2(T-t)} dt \right), \quad Y_0 = e, \quad (3)$$

where d denotes the Riemannian distance function. Note that we can always, for convenience, take the initial value to be the identity e .

Radial Process We denote by $r_v(\cdot) := d(\cdot, v)$ the radial process. Due to the radial process's singularities on $\text{Cut}(v) \cup \{v\}$, the usual Itô's formula only applies

on subsets away from the cut-locus. The extension beyond the cut-locus of a Brownian motion's radial process was due to Kendall [4]. Barden and Le [1, 5] generalized the result to M -semimartingales. The radial process of the Brownian motion (1) is given by

$$r(X_t) = r(X_0)^2 + \int_0^t \langle \nabla r(X_s), V(X_s) dB_s \rangle + \frac{1}{2} \int_0^t \Delta_G r(X_s) ds - L_s(X), \quad (4)$$

where L is the geometric local time of the cut-locus $\text{Cut}(v)$, which is non-decreasing continuous random functional increasing only when X is in $\text{Cut}(v)$ (see [1, 4, 5]). Let $W_t := \int_0^t \langle \frac{\partial}{\partial r}, V_i(X_s) \rangle dB_s^i$, which is the local-martingale part in the above equation. The quadratic variation of W_t satisfies $d[W, W]_t = dt$, by the orthonormality of $\{V_1, \dots, V_d\}$, thus W_t is a Brownian motion by Levy's characterization theorem. From the stochastic integration by parts formula and (4) the squared radial process of X satisfies

$$r(X_t)^2 = r(X_0)^2 + 2 \int_0^t r(X_s) dW_s + \int_0^t r(X_s) \Delta_G r(X_s) ds - 2 \int_0^t r(X_s) dL_s, \quad (5)$$

where dL_s is the random measure associated to $L_s(X)$.

Similarly, we obtain an expression for the squared radial process of Y . Using the shorthand notation $r_t := r_v(Y_t)$ the radial process then becomes

$$r_t^2 = r_0^2 + 2 \int_0^t r_s dW_s + \int_0^t \frac{1}{2} \Delta_G r_s^2 ds - \int_0^t \frac{r_s^2}{T-s} ds - 2 \int_0^t r_s dL_s. \quad (6)$$

Imposing a growth condition on the radial process yields an L^2 -bound on the radial process of the guided diffusion, [10]. So assume there exist constants $\nu \geq 1$ and $\lambda \in \mathbb{R}$ such that $\frac{1}{2} \Delta_G r_v^2 \leq \nu + \lambda r_v^2$ on $D \setminus \text{Cut}(v)$, for every regular domain $D \subseteq G$. Then (6) satisfies

$$\mathbb{E}[1_{t < \tau_D} r_v(Y_t)^2] \leq \left(r_v^2(e) + \nu t \left(\frac{t}{T-t} \right) \right) \left(\frac{T-t}{t} \right)^2 e^{\lambda t}, \quad (7)$$

where τ_D is the first exit time of Y from the domain D .

Girsanov Change of Measure Let B be the Brownian motion in \mathbb{R}^d defined on the filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_s), \mathbb{P})$ and X the solution of (1). The process $\frac{\nabla r_v(X_t)^2}{2(T-t)}$ is an adapted process. As X is non-explosive, we see that

$$\int_0^t \left\| \frac{\nabla r(X_s)^2}{2(T-s)} \right\|^2 ds = \int_0^t \frac{r(X_s)^2}{(T-s)^2} ds \leq C, \quad (8)$$

for every $0 \leq t < T$, almost surely, and for some fixed constant $C > 0$. Define a new measure \mathbb{Q} by

$$Z_t := \frac{d\mathbb{Q}}{d\mathbb{P}} \Big|_{\mathcal{F}_t} (X) = \exp \left\{ - \int_0^t \left\langle \frac{\nabla r(X_s)^2}{2(T-s)}, V(X_t) dB_s \right\rangle - \frac{1}{2} \int_0^t \frac{r(X_s)^2}{(T-s)^2} ds \right\}. \quad (9)$$

From (8), the process Z_t is a martingale, for $t \in [0, T)$, and \mathbb{Q}_t defines a probability measure on each \mathcal{F}_t absolutely continuous with respect to \mathbb{P} . By Girsanov's theorem (see e.g. [3, Theorem 8.1.2]) we get a new process b_s which is a Brownian motion under the probability measure \mathbb{Q} . Moreover, under the probability \mathbb{Q} , equation (1) becomes

$$dY_t = -\frac{1}{2}V_0(Y_t)dt + V_i(Y_t) \circ \left(db_t^i - \frac{r(Y_t)}{T-t} \left(\frac{\partial}{\partial r} \right)^i dt \right), \quad (10)$$

where $\left(\frac{\partial}{\partial r} \right)^i$ is the i 'th component of the unit radial vector field in the direction of v . The squared radial vector field is smooth away from $\text{Cut}(v)$ and thus we set it to zero on $\text{Cut}(v)$. Away from $\text{Cut}(v)$, the squared radial vector field is 2Log_v , which is the inverse exponential at v . The added drift term acts as a guiding term, which pulls the process towards v at time $T > 0$.

From (9), we see that $\mathbb{E}[f(Y_t)] = \mathbb{E}[f(X_t)Z_t]$. Using (5) and the identity $\Delta_G r_v = \frac{d-1}{r_v} + \frac{\partial}{\partial r_v} \log \Theta_v$ (see [9]), we equivalently write $\mathbb{E}[f(Y_t)\varphi_t] = \mathbb{E}[f(X_t)\psi_t]$, with

$$\psi_t := \exp \left\{ \frac{-r(X_t)^2}{2(T-t)} \right\} \quad \varphi_t := \exp \left\{ \int_0^t \frac{r_v(Y_s)^2}{T-s} (dA_s + dL_s) \right\}, \quad (11)$$

where $dA_s = \frac{\partial}{\partial r_v} \log \Theta_v$ is a random measure supported on $G \setminus \text{Cut}(v)$ and Θ_v is the Jacobian determinant of Exp_v .

Delyon and Hu in Lie Groups This section generalizes the result of Delyon and Hu [2, Theorem 5] to the Lie group setting. The result can be modified to incorporate a generalization of [2, Theorem 6].

Theorem 1. *Let X be the solution of (1). The SDE (3) yields a strong solution on $[0, T)$ and satisfies $\lim_{t \uparrow T} Y_t = v$ almost surely. Moreover, the conditional expectation of X given $X_T = v$ is*

$$\mathbb{E}[f(X)|X_T = v] = C \mathbb{E}[f(Y)\varphi_T], \quad (12)$$

for every \mathcal{F}_t -measurable non-negative function f on G , $t < T$, where φ_t is given in (11).

Proof. The result is a consequence of the change of measure together with Lemma 1, Lemma 2, and Lemma 3. \square

Lemma 1. *The solution of SDE (3) satisfies $\lim_{t \rightarrow T} Y_t = v$ almost surely.*

Proof. Let $\{D_n\}_{n=1}^\infty$ be an exhaustion of G , that is, the sequence consists of open, relatively compact subsets of M such that $\bar{D}_n \subseteq D_{n+1}$ and $G = \bigcup_{n=1}^\infty D_n$. Furthermore, let τ_{D_n} denote the first exit time of Y from D_n , then from (7) we have that the sequence $(\mathbb{E}[1_{\{t < \tau_{D_n}\}} r_v^2(Y_t)])_{n=1}^\infty$ is non-decreasing and bounded, hence from the monotone convergence theorem, it has a limit which is bounded

by the right-hand side of (7). Applying Jensen's inequality to the left-hand side of (7)

$$\mathbb{E}[r_v(Y_t)] \leq \left(r_v^2(e) + \nu t \left(\frac{t}{T-t} \right) \right)^{\frac{1}{2}} \left(\frac{T-t}{t} \right) e^{\frac{\lambda t}{2}}.$$

Since obviously $\mathbb{E}[r_v(Y_T)] = r_v(Y_T)\mathbb{Q}(r_v(Y_T) \neq 0)$, by Fatou's lemma $\mathbb{E}[r_v(Y_T)] \leq \liminf_{t \rightarrow T} \mathbb{E}[r(Y_t)] = 0$, we conclude that $r(Y_t) \rightarrow 0$, \mathbb{Q} -almost surely. \square

Lemma 2. *Let $0 < t_1 < t_2 < \dots < t_N < T$ and h be a continuous bounded function on G^N . With ψ_t as in (11), then*

$$\lim_{t \rightarrow T} \frac{\mathbb{E}[h(X_{t_1}, X_{t_2}, \dots, X_{t_N}) \psi_t]}{\mathbb{E}[\psi_t]} = \mathbb{E}[h(X_{t_1}, X_{t_2}, \dots, X_{t_N}) | X_T = v]. \quad (13)$$

Proof. The proof is similar to that of [2, Lemma 7]. Let (U, ϕ) be a normal chart centered at $v \in G$. First, since the cut locus of any complete connected manifold has (volume) measure zero, we can integrate indifferently in any normal chart. For any $t \in (t_N, T)$ we have

$$\mathbb{E}[h(x_{t_1}, \dots, x_{t_N}) \psi_t] = \int_G \Phi_h(t, z) e^{-\frac{r_v(z)^2}{2(T-t)}} d\text{Vol}(z) \quad (14)$$

where $d\text{Vol}(z) = \sqrt{\det(A(z))} dz$ denotes the volume measure on G , dz the Lebesgue measure, and A the metric tensor. Moreover,

$$\Phi_h(t, z) = \int_{G^N} h(z_1, \dots, z_N) p(t_1, u, z_1) \dots p(t - t_N, z_N, z) d\text{Vol}(z_1) \dots d\text{Vol}(z_N),$$

and of course $\Phi_1(t, z) = p(t, e, z)$. Using the normal chart and applying the change of variable $x = (T-t)^{1/2}y$ we get

$$(T-t)^{-\frac{d}{2}} \mathbb{E}[h(x_{t_1}, \dots, x_{t_N}) \psi_t] \xrightarrow{t \rightarrow T} \Phi_h(T, v) \det(A(v))^{\frac{d}{2}} \int_{\phi(G)} e^{-\frac{r_v(\phi^{-1}(y))^2}{2}} dy.$$

The conclusion follows from Bayes' formula. \square

Lemma 3. *With φ_t as defined above then $\varphi_t \xrightarrow{L^1} \varphi_T$.*

Proof. Note that for each $t \in [0, T)$ we have $\mathbb{E}^{\mathbb{Q}}[\varphi_t] < \infty$ as well as $\varphi_t \rightarrow \varphi_T$ almost surely by Lemma 1. The result then follows from the uniform integrability of $\{\varphi_t : t \in [0, T)\}$, which can be found in Appendix C.2 in [9]. \square

4 Importance Sampling and Metric Estimation on $\text{SO}(3)$

This section takes G to be the special orthogonal group of rotation matrices, $\text{SO}(3)$, a compact connected matrix Lie group. In the context of matrix Lie groups, computing left-invariant vector fields is straightforward.




Fig. 1: Three sample paths (a) – (c) of the guided diffusion process on $\text{SO}(3)$ visualized by its action on the basis vectors $\{e_1, e_2, e_3\}$ (red, green, blue) of \mathbb{R}^3 . The sample paths are conditioned to hit the rotation represented by the black vectors.

Numerical Simulations The Euler-Heun scheme leads to approximation of the Stratonovich integral. With a time discretization $t_1, \dots, t_k, t_k - t_{k-1} = \Delta t$ and corresponding noise $\Delta B_{t_i} \sim N(0, \Delta t)$, the numerical approximation of the Brownian motion (1) takes the form

$$x_{t_{k+1}} = x_{t_k} - \frac{1}{2} \sum_{j,i} C_{ij}^j V_i(x_{t_k}) \Delta t + \frac{v_{t_{k+1}} + V_i(v_{t_{k+1}} + x_{t_{k+1}}) \Delta B_{t_k}^i}{2} \quad (15)$$

where $v_{t_{k+1}} = V_i(x_{t_k}) \Delta B_{t_k}^i$ is only used as an intermediate value in integration. Adding the logarithmic term in (10) to (15) we obtain a numerical approximation of a guided diffusion (3). Fig. 1 shows three different sample paths from the guided diffusion conditioned to hit the rotation represented by the black vectors.

Metric Estimation on $\text{SO}(3)$ In the d -dimensional Euclidean case, importance sampling yields the estimate [7]

$$p(T, u, v) = \left(\frac{\det(A(T, v))}{2\pi T} \right)^{\frac{d}{2}} e^{-\frac{\|u-v\|_A^2}{2T}} \mathbb{E}[\varphi_T],$$


where $\|x\|_A = x^T A(0, u)x$. Thus, from the output of the importance sampling we get an estimate of the transition density. Similar to the Euclidean case, we obtain an expression for the heat kernel $p(T, e, v)$ as $p(T, e, v) = q(T, e) \mathbb{E}[\varphi_T]$, where

$$q(T, e) = \left(\frac{\det A(v)}{2\pi T} \right)^{\frac{3}{2}} \exp\left(-\frac{d(e, v)^2}{2T}\right) = \left(\frac{\det A(v)}{2\pi T} \right)^{\frac{3}{2}} \exp\left(-\frac{\|\text{Log}_v(e)\|_A^2}{2T}\right), \quad (16)$$

where the equality holds almost everywhere and $A \in \text{Sym}^+(\mathcal{G})$ denotes the metric $A(e)$. The Log_v map in (16) is the Riemannian inverse exponential map.

Figure 2 illustrate how importance sampling on $\text{SO}(3)$ leads to metric estimation of the underlying true metric, from which the Brownian motion was generated. We sampled 128 points as endpoints of a Brownian motion from the

metric $\text{diag}(0.2, 0.2, 0.8)$. We used 20 time steps and sampled 4 bridges per observation. An iterative maximum likelihood method using gradient descent with a learning rate of 0.2, and initial guess of the metric being $\text{diag}(1, 1, 1)$ yielded a convergence to the true metric. Note that in iteration the logarithmic map changes.



(a) Estimation of the unknown underlying metric using bridge sampling. Here the true metric is the diagonal matrix $\text{diag}(0.2, 0.2, 0.8)$.

(b) The iterative log-likelihood.

Fig. 2: The importance sampling technique applies to metric estimation on the Lie group $\text{SO}(3)$. Sampling a Brownian motion from an unknown underlying metric we obtain a convergence to the true underlying metric using an iterative maximum-likelihood method. Here we sampled 4 bridge processes per observation, starting from the metric $\text{diag}(1, 1, 1)$, providing a relatively smooth iterative likelihood in 2b.

References

1. Barden, D., Le, H.: Some consequences of the nature of the distance function on the cut locus in a riemannian manifold. *Journal of the LMS* (1997)
2. Delyon, B., Hu, Y.: Simulation of conditioned diffusion and application to parameter estimation. *Stochastic Processes and their Applications* (2006)
3. Hsu, E.P.: *Stochastic analysis on manifolds*, vol. 38. AMS (2002)
4. Kendall, W.S.: The radial part of brownian motion on a manifold: a semimartingale property. *The Annals of Probability* **15**(4), 1491–1500 (1987)
5. Le, H., Barden, D.: Itô correction terms for the radial parts of semimartingales on manifolds. *Probability theory and related fields* **101**(1), 133–146 (1995)
6. Liao, M.: *Lévy processes in Lie groups*, vol. 162. Cambridge university press (2004)
7. Papaspiliopoulos, O., Roberts, G.: Importance sampling techniques for estimation of diffusion models. *Statistical methods for stochastic differential equations* (2012)
8. Shigekawa, I.: Transformations of the brownian motion on a riemannian symmetric space. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* (1984)
9. Thompson, J.: Submanifold bridge processes. Ph.D. thesis, University of Warwick (2015)
10. Thompson, J.: Brownian bridges to submanifolds. *Potential Analysis* **49**(4) (2018)