

Investigating maternal brain structure and its relationship to substance use and motivational systems

Helena J.V. Rutherford^{a*}, Guido Gerig^b, Sylvain Gouttard^b, Marc N. Potenza^{a,c}, & Linda C. Mayes^a

^a Yale Child Study Center, Yale University School of Medicine, New Haven, CT, USA;

^bScientific Computing and Imaging Institute, University of Utah, Salt Lake City, UT, USA;

^c Departments of Psychiatry and Neurobiology and CASAColumbia, Yale University School of Medicine, New Haven, CT, USA

Key words: substance use, addiction, maternal brain, gray matter, behavioral inhibition / behavioral activation

*Correspondences to:

Helena J.V. Rutherford, Yale Child Study Center, Yale University, 230 South Frontage Road, New Haven, CT, 06520 USA. Phone: 203-737-3408, Fax: 203-785-7926, Email:

helena.rutherford@yale.edu

Abbreviations:

BIS – Behavioral Inhibition System

BAS - Behavioral Activation System

GM – Gray Matter

MRI – Magnetic Resonance Imaging

WM – White Matter

Acknowledgements

We thank Marion Mayes for coordinating visits and working with participants and Kara Holcomb and Laura Logan for overseeing MRI visits with participants.

This work was supported by the NIH (NIDA) grants P01 DA022446 and R01 DA026437, as well as the Anna Freud Centre (UK), the Connecticut Mental Health Center, and the National Alliance for Medical Image Computing (NA-MIC) U54 EB005149. This publication was also made possible by CTSA, Grant Number UL1 RR024139, from the National Center for Research Resources, a component of the NIH, and NIH roadmap for Medical Research. Its contents are solely the responsibility of the authors and do not necessarily represent the official view of any of the funding agencies.

The authors report that they have no financial conflicts of interest with respect to the content of this manuscript. Dr. Potenza has received financial support or compensation for the following: Dr. Potenza has consulted for and advised Somaxon, Boehringer Ingelheim, Ironwood, Lundbeck, INSYS, RiverMend Health and Shire; has received research support from the NIH,

Veteran's Administration, Mohegan Sun Casino, the National Center for Responsible Gaming, Forest Laboratories, Ortho-McNeil, Psyadon, Oy-Control/Biotie and Glaxo-SmithKline pharmaceuticals; has participated in surveys, mailings or telephone consultations related to drug addiction, impulse control disorders or other health topics; has consulted for law offices, gambling-related entities and the federal public defender's office in issues related to impulse control disorders; provides clinical care in the Connecticut Department of Mental Health and Addiction Services Problem Gambling Services Program; has performed grant reviews for the NIH and other agencies; has guest-edited journal sections; has given academic lectures in grand rounds, CME events and other clinical or scientific venues; and has generated books or book chapters for publishers of mental health texts. The other authors reported no biomedical financial interests or other conflicts of interest.

Abstract

Substance use during pregnancy and the postpartum period may have significant implications for both mother and the developing child. However, the neurobiological basis of the impact of substance use on parenting is less well understood. Here we examined the impact of maternal substance use on cortical gray matter (GM) and white matter volumes, and whether this was associated with individual differences in motivational systems of behavioral activation and inhibition. Mothers were included in the substance-using group if any addictive substance was used during pregnancy and/or in the immediate postpartum period (within 3 months of delivery). GM volume was reduced in substance-using mothers compared to non-substance-using mothers, particularly in frontal brain regions. In substance-using mothers, we also found that frontal GM was negatively correlated with levels of behavioral activation (i.e., the motivation to approach rewarding stimuli). This effect was absent in non-substance-using mothers. Taken together, these findings indicate a reduction in GM volume is associated with substance use, and that frontal GM volumetric differences may be related to approach motivation in substance-using mothers.

Introduction

Maternal substance use represents a considerable public health concern, with substance use during pregnancy and into the postpartum period continuing for many women [1]. Although substance-using mothers demonstrate difficulties during interactions with their children [2-5], the underlying neurobiological basis of this is less well understood. Converging neuroimaging studies of parents report that brain regions critical to reward, emotion and stress regulation are recruited when parents engage with infant stimuli [6-8]. In these same regions, a reduction in brain activity has been observed when substance-using mothers engage with infant stimuli [9]. This finding resonates with theoretical models that caretaking difficulties faced by substance-using mothers may reflect the dysregulation of reward and stress neurocircuitry [8, 10]. This study investigated the impact of substance use on maternal brain structure, and examined whether individual differences in motivation were associated with structural differences between substance-using and non-substance-using mothers.

Parents contribute critically to their child's development [11]; therefore, adaptation of neural architecture to facilitate parenting may have an adaptive value from evolutionary and other perspectives. While functional MRI studies have begun to interrogate the maternal brain, to our knowledge, only one previous study has measured maternal brain structure, examining gray matter (GM) volume changes during the postpartum period [12]. This study found GM volume increased from 2-4 weeks postpartum to 3-4 months postpartum in multiple regions, including prefrontal and parietal cortex. Additionally, increases in GM volume in midbrain regions were related to mothers' self-reported positive thoughts related to their babies. These findings support the potential for neurobiological reorganization at a structural level in motherhood.

Substance dependence has been associated with changes in frontal-striatal circuitries. Reductions in GM, but not white matter (WM), volume have been observed in orbitofrontal, temporal, anterior cingulate and insular regions in cocaine dependence [13]. Cocaine dependence

has also been associated with reductions in ventral striatal GM [14], and methamphetamine dependence is associated with GM reductions in the medial frontal gyrus and insula [15]. A meta-analysis reported decreased GM volume in substance-dependent participants in the prefrontal cortex [16], with GM volumes in the inferior and middle frontal gyri associated inversely with longer histories of substance use. These structural findings converge with other data illustrating that frontal cortical function is associated with multiple components of addiction [17, 18].

A recent model of parenting suggests a central role for motivation in guiding caretaking behavior in parents [8]. Therefore, understanding variability in motivational tendencies may provide insight into individual differences in caretaking in substance-using and non-substance-using parents. Converging work suggests two motivational systems underscore emotion and behavior: an approach system that drives behavior *towards* stimuli, and an avoidance system that drives behavior *away* from stimuli (e.g., [19]). These systems may map onto a behavioral activation system (BAS) and a behavioral inhibition system (BIS) that guide goal-directed behaviors [20]¹. The BAS is implicated in reward responding, guiding behavior toward desirable outcomes or stimuli. The BIS is implicated in responding to punishment, guiding behavior away from undesirable outcomes or stimuli. Notably, a recent study of non-parents evidenced BIS and BAS were associated with the neural response to infant stimuli [21] – supporting the value of examining motivational tendencies as they relate to parenting.

Carver and White (22) developed an assessment to capture variability in behavioral inhibition and activation with behavioral activation consisting of three components: (1) persistence pursuing goals (BAS-Drive); (2) engagement in seeking rewards (BAS-Fun Seeking); and (3) anticipation or response to reward receipt (BAS-Reward). The BAS is relevant to substance use given that individuals high in BAS may be more likely to seek out and have a positive response to rewards (including drugs and alcohol; [23]). Consistent with this notion, cocaine and heroin dependent participants report higher BAS scores (BAS-Drive and BAS-Fun Seeking) than do healthy control subjects [23]. Furthermore, substance use in college students positively correlated with BAS scores, specifically BAS-Fun Seeking, while only a weak correlation was found between substance use and BIS scores [24]. BAS-Drive scores have also been associated with an increased desire and intent to drink, as well as an expectation to feel relief from drinking, in participants receiving inpatient alcohol treatment [25]. Elevated scores on the BIS and all BAS subscales have also been associated with hazardous drinking in a community sample [26]. These studies suggest there may be an important coupling between substance use and motivational behavioral tendencies, particularly those relating to behavioral activation.

We examined GM and WM volumes in substance-using and non-substance-using mothers and whether structural brain differences would relate to general motivational behavioral tendencies (BIS/BAS). Given the potential damage from substance-use exposure to the developing or newborn infant, we broadly defined substance use to include any addictive substance used during pregnancy and/or postpartum. The purpose of this study was to investigate structural volumes and motivational tendencies in response to the presence (and absence) of an addictive process rather than the neurochemical effect of any one specific substance [9, 27]. We hypothesized that perinatal (i.e., during pregnancy and/or up to 3 months postpartum) substance

use would be related to differences in maternal brain structure, specifically decreased GM volume. Further, given the associations previously reported between BAS and substance use, we also hypothesized that GM volumes would be associated with BAS motivation in the substance-using mothers.

Materials and Methods

Participants

The Human Investigations Committee at Yale University School of Medicine approved all procedures, and NIDA approved a Certificate of Confidentiality for this study. Sixty-six mothers were recruited through drug treatment and rehabilitation facilities, maternity wards and posted flyers. All participants provided informed consent, and data were collected approximately 3 months (range 1-3 months) postpartum. All mothers were reimbursed \$80 and given a gift for their baby. Substance-use status was determined by self-report and urine toxicology. Women were considered substance-using ($n=31$; mean age approximately = 25.77 years; $SD = 4.89$; 9 first-time mothers) if they used any substance of abuse during pregnancy and/or within the past 30 days at time of recruitment, and/or positive toxicology screen at the time of visit. Substance-using mothers reported using only tobacco ($n=15$), tobacco and other substances ($n=10$, including marijuana, amphetamine, cocaine, heroin, alcohol, and/or other opiates), and marijuana only ($n=4$). One mother self-reported using substances but did not disclose details, and one mother was in rehabilitation. Marital status was: 26 single, 2 married, 2 divorced, and 1 mother did not report. Race/ethnicity was: 8 Caucasian, 18 African American, and 5 Hispanic/Latino.

Non-substance-using mothers ($n=35$; mean age approximately = 28.88 years; $SD = 5.70$; 27 first-time mothers) were free from tobacco or other substance use. Marital status was: 14 single and 21 married. Race/ethnicity was: 21 Caucasian, 8 African American, 2 Asian American, 2 Hispanic/Latino, and 2 mothers did not report. Consistent with evidence that there are age-related effects on brain matter volume [28], and the age difference between groups reported here, $t(62) = 2.21$, $p = .03$, age was entered where appropriate as a covariate in analyses.

Behavioral Inhibition System and Behavioral Activation System (BIS/BAS) Scale

The BIS/BAS scale [22] is a valid and reliable 24-item self-report measure designed to capture individual variability in behavioral inhibition and activation [20]. Each item is rated on a 4-point likert scale, from “1 - strongly disagree” to “4 - strongly agree”. Seven items capture behavioral inhibition, including, “If I think something unpleasant is going to happen I usually get pretty worked up”. Behavioral activation consists of three subscales: BAS-Drive (“When I want something I usually go all-out to get it”), BAS-Fun (“I’m always willing to try something new if I think it will be fun”), and BAS-Reward (“When I get something I want, I feel excited and energized”). A BAS-total score indicates the sum of all BAS subscale scores.

Image Acquisition

Magnetization prepared rapid gradient echo (MPRAGE) images (176 slices, 256 x 256 mm field of view, 256 x 256 data acquisition matrix, 2.530 s repetition time, 2.77 ms echo time, 7° flip angle, bandwidth 179 Hz/pixel) were acquired with a Siemens Trio 3T scanner (Siemens AG, Erlangen, Germany).

Automatic Tissue Segmentation

Several methods have been developed for automatic segmentation of adult brain MRI data [29-31]. Pohl et al. [32] additionally augments tissue class segmentation by a detailed parcellation of neuroanatomical structures. We used a modified version of an atlas-moderated expectation-maximization method [31]. The tool, named Atlas Based Classification (ABC), was written in ITK (Insight Consortium, 2004) and made freely available to the scientific community via the NITRC platform [33, 34]. The ABC tool takes single or multi-modal MR images as input and performs registration of a probabilistic atlas which serves as a spatial prior, bias correction, brain stripping, user-selected nonlinear filtering, and multivariate classification combined into one integrated tool. Results include tissue probability maps $p(\text{category}|x)$ for the categories of WM, GM, cerebrospinal fluid (CSF), and background (BG), and binary label maps of the maximum posterior classification, defined at each voxel location x . An additional category, the intracranial volume (ICV), is defined as the sum of WM, GM and CSF. Subdivision into lobar regions was obtained by nonlinear registration of a parcellation template to each subject's brain image, resulting in WM, GM and CSF volumes per lobe. The ABC segmentation methodology has been previously applied in large clinical studies, for example, of schizophrenia [35], and validated in a multi-site human traveling phantom study which demonstrated coefficients of variation for GM and WM in the one percent range [36].

Data Analysis

Dividing the individual structural values by the ICV for each participant was performed to normalize the data. Data from two mothers (1 substance-using; 1 non-substance-using) were excluded after boxplots of the structural data confirmed they were outliers. Analysis first focused on comparisons between total GM and WM volumes as a function of substance use. If group differences were found, the second analytic step was to examine lobe parcellation to probe the potential regional sources for substance-use differences. Greenhouse-Geisser corrections were used where applicable. The third and final analytic step was to examine associations between structural volumes where substance-use differences emerged with BIS/BAS measures. Data from the BAS subscales were not normally distributed; therefore, non-parametric analyses were used for these measures. The alpha level was defined as $p < .05$, and all data presented in figures and text are means and standard deviations.

Results

Total GM and WM Volumes

Despite statistically significant age differences between substance-use groups, this variable did not correlate with GM and WM volumes and was not included as a covariate in this analysis. Substance-using mothers presented with less total GM volume, $t(62) = 3.71$, $p < .001$, than non-substance-using mothers (Figure 1). There was no difference, $t < 1$, in total WM volume between substance-using ($M=.353$; $SD=.008$) and non-substance-using ($M=.352$; $SD=.006$) mothers.




Figure 1. Mean normalized gray matter volume as a function of substance-use group, with error bars indicating one standard deviation from the mean.

GM Parcellation

Table 1 presents the means and standard deviations for GM volume parcellation for each maternal group. To further examine GM differences, parcellated GM volume was examined using a 5 (Lobe: prefrontal, frontal, parietal, temporal, occipital) by 2 (Hemisphere: left, right) repeated-measures ANOVA with a between-group factor of substance-use status. Age was included as a covariate in the analysis after preliminary data analysis revealed age correlated with GM volume in some lobes. Age was not a significant covariate in the overall model, $F(1,61) = 2.22, p = .14$, but substance-use status was a significant between-group factor, $F(1,61) = 8.58, p < .01$. There was a main effect of lobe, $F(3,171) = 169.42, p < .001$, evidencing variability in GM volume and there being smallest GM volume in occipital and prefrontal regions (Table 1). There was a marginal interaction between lobe and substance-use status, $F(3,171) = 2.67, p = .05$. With no main effect of hemisphere, $F < 1$, or any interaction between lobe, substance-use status, and hemisphere, $F < 1$, the data were averaged across hemispheres for analysis. Independent samples t-tests showed non-substance-using mothers had more frontal cortical GM volume than substance-using mothers, $t(62) = 4.60, p < .001$. Across the other lobe regions, GM volume was comparable between the groups. Therefore, the overall reduction in total GM volume in substance-using mothers reported here seems driven by differences in GM volume in the frontal lobe.

This omnibus analysis also showed a lobe GM volume and age interaction, $F(3,171) = 7.60, p < .001$. Age did not correlate with frontal, $r(64) = -.16, p = .22$, or occipital, $r(64) = .23, p = .06$, GM volumes. There were significant inverse correlations between age and parietal GM volume, $r(64) = -.32, p = .01$, and prefrontal GM volume $r(64) = -.28, p = .02$. There was also a positive correlation between age and temporal lobe GM, $r(64) = .25, p = .04$. A lobe by hemisphere interaction was also found, $F(4,244) = 3.58, p < .01$, whereby the GM volume was larger across all lobes in the right versus left hemisphere, with the exception of the parietal lobe in which this volumetric asymmetry was reversed. There were no other statistically significant interactions between any of the remaining variables of Lobe, Hemisphere, Substance-Use Group and Age, F 's $< 3.16, p$'s $> .08$.

Please insert Table 1 about here

BIS/BAS and Frontal GM Volume

Table 2 presents BIS/BAS scores (mean and standard deviation) as a function of substance use. Although there was a non-significant trend to suggest that substance-using mothers had higher BAS-Fun scores than non-substance-using mothers, no other statistically significant differences were found between the other BAS subscales or the BIS scale as a function of substance-use group. However, we examined the relationship between BIS/BAS within each group given the statistically significant frontal GM volume differences. In substance-using-mothers, we found an inverse correlation between frontal GM volume and BAS-Fun, $r(30) = -.44, p = .02$, and BAS-Reward, $r(30) = -.39, p = .03$. There was a comparable, but not statistically significant relationship, between frontal GM volume and BAS-Drive, $r(30) = -.34, p = .06$. Figure 2 illustrates the relationship between frontal GM volume and the total BAS score, $r(30) = -.38, p = .04$. No relationship between frontal GM volume and BIS was observed, $r(30) = .07, p = .73$. We found no relationship between frontal GM volume and BIS, or any BAS subscale, in non-substance-using mothers, r 's, $< -.16, p$'s $> .36$.

Please insert Table 2 about here




Figure 2. The relationship between frontal gray matter volume and BAS-Total Score for the substance-using mothers, $r(30) = -.38, p = .04$.

Discussion

Recent work has suggested that substance use may impact maternal neural responses to infant stimuli [9]. Critically, individual differences in maternal brain structure and motivation may underlie functional correlates of substance use and infant cue perception. Past research has evidenced an important role for GM volume in maternal brain development [12]. Our finding of reduced GM volumes in substance-using mothers converges with other studies that have reported abnormalities in frontal regions associated with substance use [13, 15, 16]. While we found substance-use-related differences in overall frontal GM, understanding whether there are

regional variants in the frontal cortex will be valuable for future research. For instance, decreased GM in the orbitofrontal cortex (OFC) has been reported in substance-dependent participants [13]. The OFC contributes to reward-related processes [37, 38] and is recruited in fMRI studies where parents engage with infant stimuli [39-41]. Therefore an important extension of this work will be to relate these structural findings to maternal cognitions and behavior. While maternal behavior is likely underpinned by multiple complex neurophysiological systems [42], frontal cortical functioning may be of particular interest owing to the complexity of human parenting [43]. Our finding of structural differences in frontal GM volume will be important in guiding research questions specifically targeting the role of functions mediated by the frontal cortex in parenting. Indeed, executive functions may be associated with observable parenting behavior during parent-child interactions [44].

We investigated whether individual differences in BIS/BAS would be associated with structural brain measures. Higher levels of behavioral activation may be associated with seeking and using substances [23]. Unlike previous reports, we did not find that BAS scores differentiated substance-using from non-substance-using participants. One explanation for this null BAS finding may be that in past samples where this distinction was found, participants were typically substance-dependent [23; 25], rather than substance-using, as in the sample recruited here. However, there was a trend-level difference between groups on the BAS-Fun Seeking subscale, a measure that has previously been implicated in substance-use behaviors [24-26]. Nevertheless, owing to significant differences between groups in frontal GM volume, we assessed the relationship between structural volumes in this region and BIS/BAS. Frontal GM volumes were negatively correlated with BAS scores, specifically the BAS-Fun and BAS-Reward scales, with the correlation between GM volume and BAS-Drive not reaching statistical significance ($p = .06$). Thus, in our substance-using group, reductions in GM volume were associated with higher levels of behavioral activation (an effect absent in non-substance-using mothers). One interpretation is that decreased integrity of frontal cortical regions may associate with increased approach motivation to rewarding stimuli and events. This resonates with prior findings that impulsivity levels were negatively associated with GM volume in the left superior frontal gyrus [15]. We did not find associations between GM volume and behavioral inhibition. The role of the BIS in differentiating individuals as a function of substance use has not been consistently reported [24, 26], and the findings further suggest the value of examining behavioral activation in substance-use research.

One of the important next steps in this work will be to understand the role of BAS motivation to components of caretaking. One previous non-mother fMRI study [21] found relationships between BIS/BAS measures and neural responses to infant emotional stimuli. For instance, BAS-Drive was positively associated with activity in the right superior occipital gyrus while women viewed sad relative to neutral infant faces. A replication of this fMRI study in a maternal sample will afford the opportunity to build on the current structural findings. However, the present study adds an important component to neurobiological accounts of addiction and parenting. It has been proposed that the dysregulation through addiction of reward and stress neurocircuitry may be associated with potential difficulties many substance-using women face in caring for their children [8, 10]. Specifically, caring for infants may be relatively less rewarding and more stressful for addicted adults. Our findings suggest frontal GM reductions are associated with increased behavioral activation; therefore, approach motivation more generally may not be

compromised in these women – although the specificity of this to the caretaking role (as opposed to other activities that may interfere with parenting), as well as other social and non-social rewards, should be established.

These findings should be considered in light of limitations. There was heterogeneity in maternal substance use, without measures assessing frequency and duration of use. Although differences may exist in the effects of varying substances at a neurochemical level, the nature of addiction encompasses habitual responding underpinned by dysregulation in stress and reward systems [45, 46], consistent with a syndrome model of addiction [27]. It is also unclear when differences in GM volume emerge between substance-using and non-substance-using mothers, and whether this difference will continue across the postpartum period. A recent study reports substance-dependent individuals and their non-substance-using siblings show commonalities in brain structure and behavioral inhibition relative to unrelated control subjects – suggesting potential familial vulnerability to substance use [47]. Here our sample consisted only of mothers, and considering existing studies examining substance use and GM volume [13, 15, 16], it is likely these results may generalize to non-parent samples, although this should be empirically tested. Further, understanding what underscores differences in GM volume is critical given that this may not be related to changes in the number of neurons in GM. Finally, the maternal samples were not well matched with respect to demographics characteristics. These potential confounds represent a challenge to fully understanding the generalizability of the findings. However, with larger samples these variables may be more tightly controlled.

In summary, we found that GM volume, particularly in frontal regions, was reduced in substance-using mothers relative to non-substance-using mothers. In substance-using mothers, we also found frontal GM negatively correlated with behavioral activation. These findings add to an emerging neuroscience of human parenting and addictive behaviors, highlighting the importance of individual differences in motivational tendencies.

Footnote

¹ A fight-flight system is also recruited in the presence of threat stimuli in the immediate environment [20].

References

1. Substance Abuse and Mental Health Services Administration. Results from the 2007 National Survey on Drug Use and Health: National Findings. Rockville, MD.: 2008.
2. Johnson AL, Morrow CE, Accornero VH, Xue L, Anthony JC, Bandstra ES. Maternal Cocaine Use: Estimated Effects on Mother-Child Play Interactions in the Preschool Period. *Journal of Developmental & Behavioral Pediatrics*. 2002;23(4):191-202.
3. Molitor A, Mayes LC. Problematic dyadic interaction among toddlers and their polydrug-cocaine-using mothers. *Infant Mental Health Journal*. 2010;31(2):121-40.
4. Gottwald SR, Thurman SK. The Effects of Prenatal Cocaine Exposure on Mother--Infant Interaction and Infant Arousal in the Newborn Period. *Topics in Early Childhood Special Education*. 1994;14(2):217-31.
5. Mayes LC, Truman S. Substance abuse and parenting. In: Bornstein M, editor. *Handbook of Parenting*. Hillsdale, New Jersey: Lawrence Erlbaum; 2001. p. 329-59.
6. Swain JE. The human parental brain: In vivo neuroimaging. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2011;35(5):1242-54.
7. Rutherford HJV, Mayes LC. Primary maternal preoccupation: Using neuroimaging techniques to explore the parental brain. *Psyche*. 2011(65):973-88.
8. Rutherford HJV, Potenza MN, Mayes LC. The neurobiology of addiction and attachment. In: Suchman N, Pajulo M, Mayes LC, editors. *Parents and Substance Addiction: Developmental Approaches to Intervention*. New York: Oxford University Press; 2013.
9. Landi N, Montoya J, Kober H, Rutherford HJV, Mencl E, Worhunsky P, et al. Maternal neural responses to infant cries and faces: Relationships with substance use. *Frontiers in Psychiatry*. 2011;2(32).
10. Rutherford HJV, Williams SK, Moy S, Mayes LC, Johns JM. Disruption of maternal parenting circuitry by addictive process: rewiring of reward and stress systems. *Frontiers in Psychiatry*. 2011;2.
11. Bowlby J. *Attachment and Loss: Volume 1 Attachment*. Sydney: Pimlico; 1969.
12. Kim P, Leckman JF, Mayes LC, Feldman R, Wang X, Swain JE. The plasticity of human maternal brain: Longitudinal changes in brain anatomy during the early postpartum period. *Behavioral neuroscience*. 2010;124(5):695-700.

13. Franklin TR, Acton PD, Maldjian JA, Gray JD, Croft JR, Dackis CA, et al. Decreased gray matter concentration in the insular, orbitofrontal, cingulate, and temporal cortices of cocaine patients. *Biological Psychiatry*. 2002;51(2):134-42.
14. Barrós-Loscertales A, Garavan H, Bustamante JC, Ventura-Campos N, Llopis JJ, Belloch V, et al. Reduced striatal volume in cocaine-dependent patients. *NeuroImage*. 2011;56(3):1021-6.
15. Schwartz DL, Mitchell AD, Lahna DL, Luber HS, Huckans MS, Mitchell SH, et al. Global and local morphometric differences in recently abstinent methamphetamine-dependent individuals. *NeuroImage*. 2010;50(4):1392-401.
16. Ersche KD, Williams GB, Robbins TW, Bullmore ET. Meta-analysis of structural brain abnormalities associated with stimulant drug dependence and neuroimaging of addiction vulnerability and resilience. *Current Opinion in Neurobiology*. 2013.
17. Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *American Journal of Psychiatry*. 2002;159(10):1642-52
18. George O, Koob GF. Individual differences in prefrontal cortex function and the transition from drug use to drug dependence. *Neuroscience & Biobehavioral Reviews*. 2010;35(2):232-47.
19. Rutherford HJV, Lindell AK. Thriving and surviving: Approach and avoidance motivation and lateralization. *Emotion Review*. 2011;3(3):333-43.
20. Gray J. *The neuropsychology of anxiety: An enquiry into the functions of the septo-hippocampal system*. Oxford: Oxford University Press; 1982.
21. Montoya JL, Landi N, Kober H, Worhunsky P, Rutherford HJV, Mencl E, et al. Regional brain responses in nulliparous women to emotional infant stimuli. *PLoS biology*. 2012;7(5):e36270.
22. Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social Psychology*. 1994;67(2):319.
23. Franken I, Muris P, Georgieva I. Gray's model of personality and addiction. *Addictive Behaviors*. 2006;31(3):399-403.
24. Franken I, Muris P. BIS/BAS personality characteristics and college students' substance use. *Personality and Individual Differences*. 2006;40(7):1497-503.
25. Franken I. Behavioral approach system (BAS) sensitivity predicts alcohol craving. *Personality and Individual Differences*. 2002;32(2):349-55.

26. Hamilton KR, Sinha R, Potenza MN. Hazardous drinking and dimensions of impulsivity, behavioral approach, and inhibition in adult men and women. *Alcoholism: Clinical and Experimental Research*. 2012;36(6):434-49.
27. Shaffer HJ, LaPlante DA, LaBrie RA, Kidman RC, Donato AN, Stanton MV. Toward a syndrome model of addiction: Multiple expressions, common etiology. *Harvard Review of Psychiatry*. 2004;12(6):367-74.
28. Fjell AM, Walhovd KB. Structural brain changes in aging: courses, causes and cognitive consequences. *Reviews in the Neurosciences*. 2010;21(3):187-222.
29. Cocosco CA, Zijdenbos AP, Evans AC. A fully automatic and robust brain MRI tissue classification method. *Medical Image Analysis* 2003;7:513-27.
30. Wells WM, Kikinis R, Grimson WEL, Jolesz F. Adaptive segmentation of MRI data. *IEEE Trans Medical Imaging*. 1996;15:429–42.
31. Van Leemput K, Maes F, Vandermeulen D, Suetens P. Automated model-based tissue classification of MR images of the brain. *IEEE Trans Medical Imaging* 1999;18:897–908.
32. Pohl K, Bouix S, Nakamura M, Rohlfing T, McCarley R, Kikinis R, et al. A hierarchical algorithm for mr brain image parcellation. *IEEE Transactions on Medical Imaging* 2007;26(9):201–1212.
33. Luo XZ, Kennedy DN, Cohen Z. Neuroimaging informatics tools and resources clearinghouse (NITRC) re-source announcement. *Neuroinformatics*. 2009;7:55-6.
34. NITRC. Neuroimaging Informatics Tools and Resources Clearinghouse <http://www.nitrc.org/> 2007.
35. El-Sayed M, Steen RG, Poe MD, Bethea TC, Gerig G, Lieberman J, et al. Brain volumes in psychotic youth with schizophrenia and mood disorders. *Journal of Psychiatry and Neuroscience*. 2010;35(4):229.
36. Gouttard S, Styner M, Prastawa M, Piven J, Gerig G. Assessment of reliability of multi-site neuroimaging via traveling phantom study. *Medical Image Computing and Computer-Assisted Intervention–MICCAI 2008: Springer*; 2008. p. 263-70.
37. O'Doherty JP. Reward representations and reward-related learning in the human brain: insights from neuroimaging. *Current Opinion in Neurobiology*. 2004;14(6):769-76.
38. Gallagher M, McMahan RW, Schoenbaum G. Orbitofrontal cortex and representation of incentive value in associative learning. *Journal of Neuroscience*. 1999;19(15):6610-4.

39. Nitschke JB, Nelson EE, Rusch BD, Fox AS, Oakes TR, Davidson RJ. Orbitofrontal cortex tracks positive mood in mothers viewing pictures of their newborn infants. *NeuroImage*. 2004;21(2):583-92.
40. Noriuchi M, Kikuchi Y, Senoo A. The functional neuroanatomy of maternal love: Mother's response to infant's attachment behaviors. *Biological Psychiatry*. 2008;63(4):415-23.
41. Lorberbaum JP, Newman JD, Horwitz AR, Dubno JR, Lydiard RB, Hamner MB, et al. A potential role for thalamocingulate circuitry in human maternal behavior. *Biological Psychiatry*. 2002;51(6):431-45.
42. Gonzalez A, Atkinson L, Fleming AS. Attachment and the comparative psychobiology of mothering. In: De Haan M, Gunnar MR, editors. *Handbook of Developmental Social Neuroscience*. New York: The Guilford Press; 2009. p. 225-45.
43. Barrett J, Fleming AS. Annual Research Review: All mothers are not created equal: neural and psychobiological perspectives on mothering and the importance of individual differences. *Journal of Child Psychology and Psychiatry*. 2011;52(4):368-97.
44. Deater-Deckard K, Sewell MD, Petrill SA, Thompson LA. Maternal working memory and reactive negativity in parenting. *Psychological Science*. 2010;21(1):75-9.
45. Koob GF, Le Moal M. Drug abuse: hedonic homeostatic dysregulation. *Science (New York, NY)*. 1997;278(5335):52-8.
46. Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology*. 2009;35(1):217-38.
47. Ersche KD, Jones PS, Williams GB, Turton AJ, Robbins TW, Bullmore ET. Abnormal brain structure implicated in stimulant drug addiction. *Science (New York, NY)*. 2012;335(6068):601-4.

Table 1.

	Lobe Parcellation				
	Prefrontal	Frontal	Parietal	Temporal	Occipital
Non-SU mothers	.034 (.001)	.049 (.001)	.055 (.002)	.053 (.002)	.027 (.002)
SU mothers	.034 (.001)	.047 (.002)	.055 (.003)	.052 (.002)	.027 (.001)
<i>p</i> value	.33	< .001*	.77	.19	.22

Note. Standard deviations presented in parentheses. * Indicates statistically significant differences between groups.

Table 2.

	BIS/BAS Subscale				
	BAS-Drive	BAS-Fun	BAS-Reward	BAS-Total	BIS
Non-SU mothers	11.17 (2.59)	11.00 (1.95)	17.50 (1.88)	39.67 (5.12)	20.15 (3.62)
SU mothers	11.77 (2.61)	11.88 (2.08)	17.23 (2.16)	40.87 (5.88)	19.34 (2.80)
<i>p</i> value	.45	.06	.84	.39	.33