

Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Review Probability & Random Variables

Objective

To provide background material in support of topics in *Digital Image Processing* that are based on probability and random variables.



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Sets and Set Operations

Probability events are modeled as sets, so it is customary to begin a study of probability by defining sets and some simple operations among sets.

A *set* is a collection of objects, with each object in a set often referred to as an *element* or *member* of the set. Familiar examples include the set of all image processing books in the world, the set of prime numbers, and the set of planets circling the sun. Typically, sets are represented by uppercase letters, such as A , B , and C , and members of sets by lowercase letters, such as a , b , and c .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Sets and Set Operations (Con't)

We denote the fact that an *element* a *belongs* to set A by


$$a \in A$$

If a is not an element of A , then we write

$$a \notin A$$

A set can be specified by listing all of its elements, or by listing properties common to all elements. For example, suppose that I is the set of all integers. A set B consisting the first five nonzero integers is specified using the notation

$$B = \{1, 2, 3, 4, 5\}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Sets and Set Operations (Con't)

The set of all integers less than 10 is specified using the notation

$$C = \{c \in I \mid c < 10\}$$

which we read as "C is the set of integers such that each members of the set is less than 10." The "such that" condition is denoted by the symbol " | ". As shown in the previous two equations, the elements of the set are enclosed by curly brackets.

The set with no elements is called the *empty* or *null set*, denoted in this review by the symbol \emptyset .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Sets and Set Operations (Con't)

Two sets A and B are said to be *equal* if and only if they contain the same elements. Set equality is denoted by


$$A = B$$

If the elements of two sets are not the same, we say that the sets are *not equal*, and denote this by

$$A \neq B$$

If every element of B is also an element of A , we say that B is a *subset* of A :

$$B \subseteq A$$



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Sets and Set Operations (Con't)

Finally, we consider the concept of a *universal set*, which we denote by U and define to be the set containing all elements of interest in a given situation. For example, in an experiment of tossing a coin, there are two possible (realistic) outcomes: heads or tails. If we denote heads by H and tails by T , the universal set in this case is $\{H, T\}$. Similarly, the universal set for the experiment of throwing a single die has six possible outcomes, which normally are denoted by the face value of the die, so in this case $U = \{1, 2, 3, 4, 5, 6\}$. For obvious reasons, the universal set is frequently called the *sample space*, which we denote by S . It then follows that, for any set A , we assume that $\emptyset \subseteq A \subseteq S$, and for any element a , $a \in S$ and $a \notin \emptyset$.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Some Basic Set Operations

The operations on sets associated with basic probability theory are straightforward. The *union* of two sets A and B , denoted by

$$A \cup B$$

is the set of elements that are either in A or in B , or in both. In other words,


$$A \cup B = \{z \mid z \in A \text{ or } z \in B\}$$

Similarly, the *intersection* of sets A and B , denoted by

$$A \cap B$$

is the set of elements common to both A and B ; that is,

$$A \cap B = \{z \mid z \in A \text{ and } z \in B\}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Set Operations (Con't)

Two sets having no elements in common are said to be *disjoint* or *mutually exclusive*, in which case

$$A \cap B = \emptyset$$


The *complement* of set A is defined as

$$A^c = \{z \mid z \notin A\}$$

Clearly, $(A^c)^c = A$. Sometimes the complement of A is denoted as \overline{A} .

The *difference* of two sets A and B , denoted $A - B$, is the set of elements that belong to A , but not to B . In other words,

$$A - B = \{z \mid z \in A, z \notin B\}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Set Operations (Con't)

It is easily verified that $(A - B) = A \cap B^c$.

The union operation is applicable to multiple sets. For example the union of sets A_1, A_2, \dots, A_n is the set of points that belong to at least one of these sets. Similar comments apply to the intersection of multiple sets.

The following table summarizes several important relationships between sets. Proofs for these relationships are found in most books dealing with elementary set theory.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Set Operations (Con't)

Some Important Set Relationships

$$S^c = \emptyset; \emptyset^c = S;$$

$$A \cup A^c = S; A \cap A^c = \emptyset$$

$$A \cup \emptyset = A; A \cap \emptyset = \emptyset; S \cup \emptyset = S; S \cap \emptyset = \emptyset$$

$$A \cup A = A; A \cap A = A; A \cup S = S; A \cap S = A$$


$$A \cup B = B \cup A; A \cap B = B \cap A$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$(A \cup B) \cup C = A \cup (B \cup C) = A \cup B \cup C$$

$$(A \cap B) \cap C = A \cap (B \cap C) = A \cap B \cap C$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Set Operations (Con't)

It often is quite useful to represent sets and sets operations in a so-called *Venn diagram*, in which S is represented as a rectangle, sets are represented as areas (typically circles), and points are associated with elements. The following example shows various uses of Venn diagrams.

Example: The following figure shows various examples of Venn diagrams. The shaded areas are the result (sets of points) of the operations indicated in the figure. The diagrams in the top row are self explanatory. The diagrams in the bottom row are used to prove the validity of the expression

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C) - A \cap B \cap C$$

which is used in the proof of some probability relationships.




Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Set Operations (Con't)





Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Probability

A *random experiment* is an experiment in which it is not possible to predict the outcome. Perhaps the best known random experiment is the tossing of a coin. Assuming that the coin is not biased, we are used to the concept that, on average, half the tosses will produce heads (H) and the others will produce tails (T). This is intuitive and we do not question it. In fact, few of us have taken the time to verify that this is true. If we did, we would make use of the concept of relative frequency. Let n denote the total number of tosses, n_H the number of heads that turn up, and n_T the number of tails. Clearly,

$$n_H + n_T = n.$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Dividing both sides by n gives

$$\frac{n_H}{n} + \frac{n_T}{n} = 1.$$

The term n_H/n is called the *relative frequency* of the event we have denoted by H , and similarly for n_T/n . If we performed the tossing experiment a large number of times, we would find that each of these relative frequencies tends toward a stable, limiting value. We call this value the *probability of the event*, and denoted it by $P(\text{event})$.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


In the current discussion the probabilities of interest are $P(H)$ and $P(T)$. We know in this case that $P(H) = P(T) = 1/2$. Note that the event of an experiment need not signify a single outcome. For example, in the tossing experiment we could let D denote the event "heads or tails," (note that the event is now a set) and the event E , "neither heads nor tails." Then, $P(D) = 1$ and $P(E) = 0$.

The first important property of P is that, for an event A ,

$$0 \leq P(A) \leq 1.$$

That is, the probability of an event is a positive number bounded by 0 and 1. For the certain event, S ,

$$P(S) = 1.$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Here the certain event means that the outcome is from the universal or sample set, S . Similarly, we have that for the impossible event, S^c

$$P(S^c) = 0.$$

This is the probability of an event being outside the sample set. In the example given at the end of the previous paragraph, $S = D$ and $S^c = E$.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

The *event* that either events A *or* B *or both* have occurred is simply the union of A and B (recall that events can be sets). Earlier, we denoted the union of two sets by $A \cup B$. One often finds the equivalent notation $A+B$ used interchangeably in discussions on probability. Similarly, the event that *both* A *and* B occurred is given by the intersection of A and B , which we denoted earlier by $A \cap B$. The equivalent notation AB is used much more frequently to denote the occurrence of both events in an experiment.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Relative Frequency & Prob. (Con't)

Suppose that we conduct our experiment n times. Let n_1 be the number of times that only event A occurs; n_2 the number of times that B occurs; n_3 the number of times that AB occurs; and n_4 the number of times that neither A nor B occur. Clearly, $n_1 + n_2 + n_3 + n_4 = n$. Using these numbers we obtain the following relative frequencies:

$$\frac{n_A}{n} = \frac{n_1 + n_3}{n}$$

$$\frac{n_B}{n} = \frac{n_2 + n_3}{n}$$

$$\frac{n_{AB}}{n} = \frac{n_3}{n}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


and

$$\begin{aligned}\frac{n_{A \cup B}}{n} &= \frac{n_1 + n_2 + n_3}{n} \\ &= \frac{(n_1 + n_3) + (n_2 + n_3) - n_3}{n} \\ &= \frac{n_A}{n} + \frac{n_B}{n} - \frac{n_{AB}}{n}.\end{aligned}$$

Using the previous definition of probability based on relative frequencies we have the important result

$$P(A \cup B) = P(A) + P(B) - P(AB).$$

If A and B are *mutually exclusive* it follows that the set AB is empty and, consequently, $P(AB) = 0$.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

The relative frequency of event A occurring, *given that* event B has occurred, is given by

$$\begin{aligned}\frac{n_{A/B}}{n} &= \frac{\frac{n_{AB}}{n}}{\frac{n_B}{n}} \\ &= \frac{n_3}{n_2 + n_3}.\end{aligned}$$

This *conditional probability* is denoted by $P(A/B)$, where we note the use of the symbol “/” to denote conditional occurrence. It is common terminology to refer to $P(A/B)$ as the *probability of A given B* .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Similarly, the relative frequency of B occurring, given that A has occurred is

$$\begin{aligned}\frac{n_{B/A}}{n} &= \frac{\frac{n_{AB}}{n}}{\frac{n_A}{n}} \\ &= \frac{n_3}{n_1 + n_3}.\end{aligned}$$

We call this relative frequency *the probability of B given A* , and denote it by $P(B/A)$.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

A little manipulation of the preceding results yields the following important relationships

$$P(A/B) = \frac{P(B/A)P(A)}{P(B)}$$


and

$$P(AB) = P(A)P(B/A) = P(B)P(A/B).$$

The second expression may be written as

$$P(B/A) = \frac{P(A/B)P(B)}{P(A)}$$

which is known as *Bayes' theorem*, so named after the 18th century mathematician Thomas Bayes.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Example: Suppose that we want to extend the expression

$$P(A \cup B) = P(A) + P(B) - P(AB)$$

to three variables, A , B , and C . Recalling that AB is the same as $A \cap B$, we replace B by $B \cup C$ in the preceding equation to obtain


$$P(A \cup B \cup C) = P(A) + P(B \cup C) - P(A \cap [B \cup C]).$$

The second term in the right can be written as

$$P(B \cup C) = P(B) + P(C) - P(BC).$$

From the Table discussed earlier, we know that

$$A \cap [B \cup C] = (A \cap B) \cup (A \cap C)$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

SO,

$$\begin{aligned}P(A \cap [B \cup C]) &= P([A \cap B] \cup [A \cap C]) \\ &= P(AB \cup AC) \\ &= P(AB) + P(AC) - P(ABC).\end{aligned}$$


Collecting terms gives us the final result

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(AB) - P(AC) - P(BC) + P(ABC).$$

Proceeding in a similar fashion gives

$$P(ABC) = P(A)P(B/A)P(C/AB).$$

The preceding approach can be used to generalize these expressions to N events.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


If A and B are *statistically independent*, then $P(B/A) = P(B)$ and it follows that

$$\begin{aligned}P(A/B) &= P(A) \\ P(B/A) &= P(B)\end{aligned}$$

and

$$P(AB) = P(A)P(B).$$

It was stated earlier that if sets (events) A and B are *mutually exclusive*, then $A \cap B = \emptyset$ from which it follows that $P(AB) = P(A \cap B) = 0$. As was just shown, the two sets are statistically independent if $P(AB) = P(A)P(B)$, which we assume to be nonzero in general. *Thus, we conclude that for two events to be statistically independent, they cannot be mutually exclusive.*



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

For three events A , B , and C to be independent, it must be true that


$$P(AB) = P(A)P(B)$$

$$P(AC) = P(A)P(C)$$

$$P(BC) = P(B)P(C)$$

and

$$P(ABC) = P(A)P(B)P(C).$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


In general, for N events to be statistically independent, it must be true that, for all combinations $1 \leq i \leq j \leq k \leq \dots \leq N$

$$P(A_i A_j) = P(A_i)P(A_j)$$

$$P(A_i A_j A_k) = P(A_i)P(A_j)P(A_k)$$

$$\vdots$$

$$P(A_1 A_2 \cdots A_N) = P(A_1)P(A_2) \cdots P(A_N).$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Example: (a) An experiment consists of throwing a single die twice. The probability of any of the six faces, 1 through 6, coming up in either experiment is $1/6$. Suppose that we want to find the probability that a 2 comes up, followed by a 4. These two events are statistically independent (the second event does not depend on the outcome of the first). Thus, letting A represent a 2 and B a 4,

$$P(AB) = P(A)P(B) = \frac{1}{6} \times \frac{1}{6} = \frac{1}{36}.$$

We would have arrived at the same result by defining "2 followed by 4" to be a single event, say C . The sample set of all possible outcomes of two throws of a die is 36. Then, $P(C)=1/36$.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


Example (Con't): (b) Consider now an experiment in which we draw one card from a standard card deck of 52 cards. Let A denote the event that a king is drawn, B denote the event that a queen or jack is drawn, and C the event that a diamond-face card is drawn. A brief review of the previous discussion on relative frequencies would show that

$$P(A) = \frac{4}{52},$$

$$P(B) = \frac{8}{52},$$

and

$$P(C) = \frac{13}{52}.$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)


Example (Con't): Furthermore,

$$P(AC) = P(A \cap C) = P(A)P(C) = \frac{1}{52}$$

and

$$P(BC) = P(B \cap C) = P(B)P(C) = \frac{2}{52}.$$

Events A and B are mutually exclusive (we are drawing only one card, so it would be impossible to draw a king and a queen or jack simultaneously). Thus, it follows from the preceding discussion that $P(AB) = P(A \cap B) = 0$ [and also that $P(AB) \neq P(A)P(B)$].



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Example (Con't): (c) As a final experiment, consider the deck of 52 cards again, and let A_1 , A_2 , A_3 , and A_4 represent the events of drawing an ace in each of four successive draws. If we replace the card drawn before drawing the next card, then the events are statistically independent and it follows that

$$\begin{aligned} P(A_1A_2A_3A_4) &= P(A_1)P(A_2)P(A_3)P(A_4) \\ &= \left[\frac{4}{52} \right]^4 \approx 3.5 \times 10^{-5}. \end{aligned}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Relative Frequency & Prob. (Con't)

Example (Con't): Suppose now that we do not replace the cards that are drawn. The events then are no longer statistically independent. With reference to the results in the previous example, we write

$$\begin{aligned}P(A_1A_2A_3A_4) &= P(A_1)P(A_2A_3A_4/A_1) \\ &= P(A_1)P(A_2/A_1)P(A_3A_4/A_1A_2) \\ &= P(A_1)P(A_2/A_1)P(A_3/A_1A_2)P(A_4/A_1A_2A_3) \\ &= \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49} \approx 3.7 \times 10^{-6}.\end{aligned}$$

Thus we see that not replacing the drawn card reduced our chances of drawing four successive aces by a factor of close to 10. This significant difference is perhaps larger than might be expected from intuition.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables

Random variables often are a source of confusion when first encountered. This need not be so, as the concept of a random variable is in principle quite simple. A *random variable*, x , is a real-valued function *defined* on the events of the sample space, S . In words, for each event in S , there is a real number that is the corresponding value of the random variable. Viewed yet another way, a random variable maps each event in S onto the real line. That is it. A simple, straightforward definition.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

Part of the confusion often found in connection with random variables is the fact that they are *functions*. The notation also is partly responsible for the problem. In other words, although typically the notation used to denote a random variable is as we have shown it here, x , or some other appropriate variable, to be strictly formal, a random variable should be written as a function $x(\cdot)$ where the argument is a specific event being considered. However, this is seldom done, and, in our experience, trying to be formal by using function notation complicates the issue more than the clarity it introduces. Thus, we will opt for the less formal notation, with the warning that it must be kept clearly in mind that random variables are functions.



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

Example: Consider again the experiment of drawing a single card from a standard deck of 52 cards. Suppose that we define the following events. A : a heart; B : a spade; C : a club; and D : a diamond, so that $S = \{A, B, C, D\}$. A random variable is easily defined by letting $x = 1$ represent event A , $x = 2$ represent event B , and so on.

As a second illustration, consider the experiment of throwing a single die and observing the value of the up-face. We can define a random variable as the numerical outcome of the experiment (i.e., 1 through 6), but there are many other possibilities. For example, a binary random variable could be defined simply by letting $x = 0$ represent the event that the outcome of throw is an even number and $x = 1$ otherwise.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

Note the important fact in the examples just given that the probability of the events have not changed; all a random variable does is map events onto the real line.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

Thus far we have been concerned with random variables whose values are discrete. To handle *continuous random variables* we need some additional tools. In the discrete case, the probabilities of events are numbers between 0 and 1. When dealing with continuous quantities (which are not denumerable) we can no longer talk about the "probability of an event" because that probability is zero. This is not as unfamiliar as it may seem. For example, given a continuous function we know that the area of the function between two limits a and b is the integral from a to b of the function. However, the area *at a point* is zero because the integral from, say, a to a is zero. We are dealing with the same concept in the case of continuous random variables.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Random Variables (Con't)

Thus, instead of talking about the probability of a specific value, we talk about the probability that the value of the random variable lies in a specified *range*. In particular, we are interested in the probability that the random variable is less than or equal to (or, similarly, greater than or equal to) a specified constant a . We write this as

$$F(a) = P(x \leq a).$$

If this function is given for all values of a (i.e., $-\infty < a < \infty$), then the values of random variable x have been defined. Function F is called the *cumulative probability distribution function* or simply the *cumulative distribution function* (cdf). The shortened term *distribution function* also is used.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Random Variables (Con't)

Observe that the notation we have used makes no distinction between a random variable and the values it assumes. If confusion is likely to arise, we can use more formal notation in which we let capital letters denote the random variable and lowercase letters denote its values. For example, the cdf using this notation is written as

$$F_X(x) = P(X \leq x).$$

When confusion is not likely, the cdf often is written simply as $F(x)$. This notation will be used in the following discussion when speaking generally about the cdf of a random variable.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Random Variables (Con't)

Due to the fact that it is a probability, the cdf has the following properties:

1. $F(-\infty) = 0$
2. $F(\infty) = 1$
3. $0 \leq F(x) \leq 1$
4. $F(x_1) \leq F(x_2)$ if $x_1 < x_2$
5. $P(x_1 < x \leq x_2) = F(x_2) - F(x_1)$
6. $F(x^+) = F(x)$,

where $x^+ = x + \varepsilon$, with ε being a positive, infinitesimally small number.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Random Variables (Con't)

The *probability density function* (pdf) of random variable x is defined as the derivative of the cdf:

$$p(x) = \frac{dF(x)}{dx}.$$

The term *density function* is commonly used also. The pdf satisfies the following properties:

1. $p(x) \geq 0$ for all x
2. $\int_{-\infty}^{\infty} p(x)dx = 1$
3. $F(x) = \int_{-\infty}^x p(a)da$, where a is a dummy variable
4. $P(x_1 < x \leq x_2) = \int_{x_1}^{x_2} p(x)dx$.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

The preceding concepts are applicable to discrete random variables. In this case, there is a finite no. of events and we talk about *probabilities*, rather than probability density functions. Integrals are replaced by summations and, sometimes, the random variables are subscripted. For example, in the case of a discrete variable with N possible values we would denote the probabilities by $P(x_i)$, $i=1, 2, \dots, N$.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Random Variables (Con't)

In Sec. 3.3 of the book we used the notation $p(r_k)$, $k = 0, 1, \dots, L - 1$, to denote the *histogram* of an image with L possible gray levels, r_k , $k = 0, 1, \dots, L - 1$, where $p(r_k)$ is the probability of the k th gray level (random event) occurring. The discrete random variables in this case are gray levels. It generally is clear from the context whether one is working with continuous or discrete random variables, and whether the use of subscripting is necessary for clarity. Also, uppercase letters (e.g., P) are frequently used to distinguish between probabilities and probability density functions (e.g., p) when they are used together in the same discussion.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Random Variables (Con't)

If a random variable x is *transformed* by a monotonic transformation function $T(x)$ to produce a new random variable y , the probability density function of y can be obtained from knowledge of $T(x)$ and the probability density function of x , as follows:

$$p_y(y) = p_x(x) \left| \frac{dx}{dy} \right|$$

where the subscripts on the p 's are used to denote the fact that they are different functions, and the vertical bars signify the absolute value. A function $T(x)$ is *monotonically increasing* if $T(x_1) < T(x_2)$ for $x_1 < x_2$, and *monotonically decreasing* if $T(x_1) > T(x_2)$ for $x_1 < x_2$. The preceding equation is valid if $T(x)$ is an increasing or decreasing monotonic function.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Expected Value and Moments

The *expected value* of a function $g(x)$ of a *continuous* random variable is defined as

$$E[g(x)] = \int_{-\infty}^{\infty} g(x)p(x)dx.$$

If the random variable is *discrete* the definition becomes

$$E[g(x)] = \sum_{i=1}^N g(x_i)P(x_i).$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)


The expected value is one of the operations used most frequently when working with random variables. For example, the expected value of random variable x is obtained by letting $g(x) = x$:

$$E[x] = \bar{x} = m = \int_{-\infty}^{\infty} xp(x)dx$$

when x is continuous and

$$E[x] = \bar{x} = m = \sum_{i=1}^N x_i P(x_i)$$

when x is discrete. The expected value of x is equal to its **average** (or **mean**) **value**, hence the use of the equivalent notation \bar{x} and m .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)


The *variance* of a random variable, denoted by σ^2 , is obtained by letting $g(x) = x^2$ which gives

$$\sigma^2 = E[x^2] = \int_{-\infty}^{\infty} x^2 p(x) dx$$

for continuous random variables and

$$\sigma^2 = E[x^2] = \sum_{i=1}^N x_i^2 P(x_i)$$

for discrete variables.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)


Of particular importance is the variance of random variables that have been *normalized* by subtracting their mean. In this case, the variance is

$$\sigma^2 = E[(x - m)^2] = \int_{-\infty}^{\infty} (x - m)^2 p(x) dx$$

and

$$\sigma^2 = E[(x - m)^2] = \sum_{i=1}^N (x_i - m)^2 P(x_i)$$

for continuous and discrete random variables, respectively. The square root of the variance is called the *standard deviation*, and is denoted by σ .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)


We can continue along this line of thought and define the n th *central moment* of a continuous random variable by letting $g(x) = (x - m)^n$:

$$\mu_n = E[(x - m)^n] = \int_{-\infty}^{\infty} (x - m)^n p(x) dx$$

and

$$\mu_n = E[(x - m)^n] = \sum_{i=1}^N (x_i - m)^n P(x_i)$$

for discrete variables, where we assume that $n \geq 0$. Clearly, $\mu_0=1$, $\mu_1=0$, and $\mu_2=\sigma^2$. The term *central* when referring to moments indicates that the mean of the random variables has been subtracted out. The moments defined above in which the mean is not subtracted out sometimes are called *moments about the origin*.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)

In image processing, moments are used for a variety of purposes, including histogram processing, segmentation, and description. In general, moments are used to characterize the probability density function of a random variable. For example, the second, third, and fourth central moments are intimately related to the *shape* of the probability density function of a random variable. The second central moment (the centralized variance) is a measure of *spread* of values of a random variable about its mean value, the third central moment is a measure of *skewness* (bias to the left or right) of the values of x about the mean value, and the fourth moment is a relative measure of *flatness*. In general, knowing all the moments of a density specifies that density.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)

Example: Consider an experiment consisting of repeatedly firing a rifle at a target, and suppose that we wish to characterize the behavior of bullet impacts on the target in terms of whether we are shooting high or low.. We divide the target into an upper and lower region by passing a horizontal line through the bull's-eye. The events of interest are the vertical distances from the center of an impact hole to the horizontal line just described. Distances above the line are considered positive and distances below the line are considered negative. The distance is zero when a bullet hits the line.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Expected Value & Moments (Con't)

In this case, we define a random variable directly as the value of the distances in our sample set. Computing the mean of the random variable indicates whether, *on average*, we are shooting high or low. If the mean is zero, we know that the average of our shots are on the line. However, the mean does not tell us how far our shots deviated from the horizontal. The variance (or standard deviation) will give us an idea of the *spread of the shots*. A small variance indicates a tight grouping (with respect to the mean, and in the vertical position); a large variance indicates the opposite. Finally, a third moment of zero would tell us that the spread of the shots is symmetric about the mean value, a positive third moment would indicate a high bias, and a negative third moment would tell us that we are shooting low more than we are shooting high with respect to the mean location.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


The Gaussian Probability Density Function

Because of its importance, we will focus in this tutorial on the *Gaussian probability density function* to illustrate many of the preceding concepts, and also as the basis for generalization to more than one random variable. The reader is referred to Section 5.2.2 of the book for examples of other density functions.

A random variable is called *Gaussian* if it has a probability density of the form

$$p(x) = \frac{1}{\sqrt{2\pi} \sigma} e^{-(x-m)^2/\sigma^2}$$

where m and σ are as defined in the previous section. The term *normal* also is used to refer to the Gaussian density. A plot and properties of this density function are given in Section 5.2.2 of the book.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

The Gaussian PDF (Con't)

The cumulative distribution function corresponding to the Gaussian density is

$$\begin{aligned} F(x) &= \int_{-\infty}^x p(x) dx \\ &= \frac{1}{\sqrt{2\pi} \sigma} \int_{-\infty}^x e^{-(x-m)^2/\sigma^2} dx. \end{aligned}$$

which, as before, we interpret as the probability that the random variable lies between minus infinite and an arbitrary value x . This integral has no known closed-form solution, and it must be solved by numerical or other approximation methods. Extensive tables exist for the Gaussian cdf.



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Several Random Variables

In the previous example, we used a single random variable to describe the behavior of rifle shots with respect to a horizontal line passing through the bull's-eye in the target. Although this is useful information, it certainly leaves a lot to be desired in terms of telling us how well we are shooting with respect to the center of the target. In order to do this we need two random variables that will map our events onto the xy -plane. It is not difficult to see how if we wanted to describe events in 3-D space we would need three random variables. In general, we consider in this section the case of n random variables, which we denote by x_1, x_2, \dots, x_n (the use of n here is not related to our use of the same symbol to denote the n th moment of a random variable).



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


Several Random Variables (Con't)

It is convenient to use vector notation when dealing with several random variables. Thus, we represent a *vector random variable* \mathbf{x} as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}.$$

Then, for example, the cumulative distribution function introduced earlier becomes

$$\begin{aligned} F(\mathbf{a}) &= F(a_1, a_2, \dots, a_n) \\ &= P\{x_1 \leq a_1, x_2 \leq a_2, \dots, x_n \leq a_n\} \end{aligned}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Several Random Variables (Con't)

when using vectors. As before, when confusion is not likely, the *cdf of a random variable vector* often is written simply as $F(\mathbf{x})$. This notation will be used in the following discussion when speaking generally about the cdf of a random variable vector.

As in the single variable case, the *probability density function of a random variable vector* is defined in terms of derivatives of the cdf; that is,

$$\begin{aligned} p(\mathbf{x}) &= p(x_1, x_2, \dots, x_n) \\ &= \frac{\partial^n F(x_1, x_2, \dots, x_n)}{\partial x_1 \partial x_2 \cdots \partial x_n}. \end{aligned}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Several Random Variables (Con't)

The *expected value* of a function of \mathbf{x} is defined basically as before:

$$\begin{aligned} E[g(\mathbf{x})] &= E[g(x_1, x_2, \dots, x_n)] \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \cdots \int_{-\infty}^{\infty} g(x_1, x_2, \dots, x_n) p(x_1, x_2, \dots, x_n) dx_1 dx_2 \cdots dx_n. \end{aligned}$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Several Random Variables (Con't)

Cases dealing with expectation operations involving pairs of elements of \mathbf{x} are particularly important. For example, the joint moment (about the origin) of order kq between variables x_i and x_j

$$\eta_{kq}(i,j) = E[x_i^k x_j^q] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x_i^k x_j^q p(x_i, x_j) dx_i dx_j.$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Several Random Variables (Con't)

When working with any two random variables (any two elements of \mathbf{x}) it is common practice to simplify the notation by using x and y to denote the random variables. In this case the joint moment just defined becomes

$$\eta_{kq} = E[x^k y^q] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x^k y^q p(x, y) dx dy.$$

It is easy to see that η_{k0} is the k th moment of x and η_{0q} is the q th moment of y , as defined earlier.



Digital Image Processing, 3rd ed.

Gonzalez & Woods


Review of Probability

www.ImageProcessingPlace.com

Several Random Variables (Con't)

The moment $\eta_{11} = E[xy]$ is called the **correlation** of x and y . As discussed in Chapters 4 and 12 of the book, correlation is an important concept in image processing. In fact, it is important in most areas of signal processing, where typically it is given a special symbol, such as R_{xy} :

$$R_{xy} = \eta_{11} = E[xy] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xyp(x,y)dxdy.$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Several Random Variables (Con't)


If the condition

$$R_{xy} = E[x]E[y]$$

holds, then the two random variables are said to be *uncorrelated*. From our earlier discussion, we know that if x and y are *statistically independent*, then $p(x, y) = p(x)p(y)$, in which case we write

$$R_{xy} = \int_{-\infty}^{\infty} xp(x)dx \int_{-\infty}^{\infty} yp(y)dy = E[x]E[y].$$

Thus, we see that *if two random variables are statistically independent then they are also uncorrelated*. The converse of this statement is *not* true in general.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Several Random Variables (Con't)


The joint central moment of order kq involving random variables x and y is defined as

$$\begin{aligned}\mu_{kq} &= E[(x - m_x)^k (y - m_y)^q] \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - m_x)^k (y - m_y)^q p(x, y) dx dy\end{aligned}$$

where $m_x = E[x]$ and $m_y = E[y]$ are the means of x and y , as defined earlier. We note that

$$\mu_{20} = E[(x - m_x)^2] \quad \text{and} \quad \mu_{02} = E[(y - m_y)^2]$$

are the variances of x and y , respectively.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Several Random Variables (Con't)

The moment μ_{11}

$$\begin{aligned}\mu_{11} &= E[(x - m_x)(y - m_y)] \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x - m_x)(y - m_y)p(x, y) dx dy\end{aligned}$$

is called the *covariance* of x and y . As in the case of correlation, the covariance is an important concept, usually given a special symbol such as C_{xy} .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Several Random Variables (Con't)

By direct expansion of the terms inside the expected value brackets, and recalling the $m_x = E[x]$ and $m_y = E[y]$, it is straightforward to show that

$$\begin{aligned}C_{xy} &= E[xy] - m_y E[x] - m_x E[y] + m_x m_y \\ &= E[xy] - E[x]E[y] \\ &= R_{xy} - E[x]E[y].\end{aligned}$$

From our discussion on correlation, we see that the covariance is zero if the random variables are either uncorrelated *or* statistically independent. This is an important result worth remembering.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

Several Random Variables (Con't)

If we divide the covariance by the square root of the product of the variances we obtain

$$\begin{aligned}\gamma &= \frac{\mu_{11}}{\sqrt{\mu_{20}\mu_{02}}} \\ &= \frac{C_{xy}}{\sigma_x\sigma_y} \\ &= E\left[\frac{(x - m_x)}{\sigma_x} \frac{(y - m_y)}{\sigma_y}\right].\end{aligned}$$

The quantity γ is called the *correlation coefficient* of random variables x and y . It can be shown that γ is in the range $-1 \leq \gamma \leq 1$ (see Problem 12.5). As discussed in Section 12.2.1, the correlation coefficient is used in image processing for matching.



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

The Multivariate Gaussian Density

As an illustration of a probability density function of more than one random variable, we consider the *multivariate Gaussian probability density function*, defined as

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} |\mathbf{C}|^{1/2}} e^{-\frac{1}{2} [(\mathbf{x}-\mathbf{m})^T \mathbf{C}^{-1} (\mathbf{x}-\mathbf{m})]}$$

where n is the *dimensionality* (number of components) of the random vector \mathbf{x} , \mathbf{C} is the *covariance matrix* (to be defined below), $|\mathbf{C}|$ is the determinant of matrix \mathbf{C} , \mathbf{m} is the *mean vector* (also to be defined below) and T indicates transposition (see the review of matrices and vectors).



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com


The Multivariate Gaussian Density (Con't)

The *mean vector* is defined as

$$\mathbf{m} = E[\mathbf{x}] = \begin{bmatrix} E[x_1] \\ E[x_2] \\ \vdots \\ E[x_n] \end{bmatrix}$$

and the *covariance matrix* is defined as

$$\mathbf{C} = E[(\mathbf{x} - \mathbf{m})(\mathbf{x} - \mathbf{m})^T].$$



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability


www.ImageProcessingPlace.com

The Multivariate Gaussian Density (Con't)

The element of \mathbf{C} are the covariances of the elements of \mathbf{x} , such that

$$c_{ij} = C_{x_i x_j} = E[(x_i - m_i)(x_j - m_j)]$$

where, for example, x_i is the i th component of \mathbf{x} and m_i is the i th component of \mathbf{m} .



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

The Multivariate Gaussian Density (Con't)

Covariance matrices are *real* and *symmetric* (see the review of matrices and vectors). The elements along the main diagonal of \mathbf{C} are the variances of the elements \mathbf{x} , such that $c_{ii} = \sigma_{x_i}^2$. When all the elements of \mathbf{x} are uncorrelated or statistically independent, $c_{ij} = 0$, and the covariance matrix becomes a *diagonal matrix*. If all the variances are equal, then the covariance matrix becomes proportional to the *identity matrix*, with the constant of proportionality being the variance of the elements of \mathbf{x} .



Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

The Multivariate Gaussian Density (Con't)

Example: Consider the following *bivariate* ($n = 2$) Gaussian probability density function


$$p(\mathbf{x}) = \frac{1}{(2\pi)^{n/2} |\mathbf{C}|^{1/2}} e^{-\frac{1}{2} [(\mathbf{x}-\mathbf{m})^T \mathbf{C}^{-1} (\mathbf{x}-\mathbf{m})]}$$

with

$$\mathbf{m} = \begin{bmatrix} m_1 \\ m_2 \end{bmatrix}$$

and

$$\mathbf{C} = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$$



Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

The Multivariate Gaussian Density (Con't)

where, because \mathbf{C} is known to be symmetric, $c_{12} = c_{21}$. A schematic diagram of this density is shown in Part (a) of the following figure. Part (b) is a horizontal slice of Part (a). From the review of vectors and matrices, we know that the main directions of data spread are in the directions of the eigenvectors of \mathbf{C} . Furthermore, if the variables are uncorrelated or statistically independent, the covariance matrix will be diagonal and the eigenvectors will be in the same direction as the coordinate axes x_1 and x_2 (and the ellipse shown would be oriented along the x_1 - and x_2 -axis). If, the variances along the main diagonal are equal, the density would be symmetrical in all directions (in the form of a bell) and Part (b) would be a circle. Note in Parts (a) and (b) that the density is centered at the mean values (m_1, m_2) .





Digital Image Processing, 3rd ed.


Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

The Multivariate Gaussian Density (Con't)





Digital Image Processing, 3rd ed.

Gonzalez & Woods

Review of Probability

www.ImageProcessingPlace.com

Linear Transformations of Random Variables

As discussed in the *Review of Matrices and Vectors*, a linear transformation of a vector \mathbf{x} to produce a vector \mathbf{y} is of the form $\mathbf{y} = \mathbf{A}\mathbf{x}$. Of particular importance in our work is the case when the rows of \mathbf{A} are the eigenvectors of the covariance matrix. Because \mathbf{C} is real and symmetric, we know from the discussion in the *Review of Matrices and Vectors* that it is always possible to find n orthonormal eigenvectors from which to form \mathbf{A} . The implications of this are discussed in considerable detail at the end of the *Review of Matrices and Vectors*, which we recommend should be read again as a conclusion to the present discussion.