In all that follows, a, b, c, c_1 and c_2 are arbitrary real constants, and f and g are functions.

n is any positive integer.

r is any rational number.

The notation \pm means "plus or minus", and \mp is "minus or plus". When used in the same equations, these symbols retain their ordering. E.g., $\pm x = \mp 3$ means both "+x = -3" and "-x = +3".

Trigonometric identities and alues:

$$\sin (a \pm b) = \sin a \cos b \pm \cos a \sin b$$

$$\sin (2a) = 2 \sin a \cos a$$

$$\sin^2 a + \cos^2 a = 1$$

$$\sin 0 = \cos \frac{\pi}{2} = 0$$

$$\sin \frac{\pi}{6} = \cos \frac{\pi}{3} = \frac{1}{2}$$

$$\sin \frac{\pi}{4} = \cos \frac{\pi}{4} = \frac{1}{\sqrt{2}}$$

$$\cos (a \pm b) = \cos a \cos b \mp \sin a \sin b$$

$$\cos (2a) = \cos^2 a - \sin^2 a$$

$$\sin \frac{\pi}{2} = \cos 0 = 1$$

$$\sin \frac{\pi}{3} = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}$$

Limits and continuity:

$$\lim_{x \to c} c_1 = c_1 \qquad \qquad \lim_{x \to c} x = c$$

$$\lim_{x \to c} (c_1 f(x) \pm c_2 g(x)) = c_1 \lim_{x \to c} f(x) \pm c_2 \lim_{x \to c} g(x) \qquad \qquad \lim_{x \to c} f(x) g(x) = \left(\lim_{x \to c} f(x)\right) \left(\lim_{x \to c} g(x)\right)$$

$$\lim_{x \to c} \frac{f(x)}{g(x)} = \frac{\lim_{x \to c} f(x)}{\lim_{x \to c} g(x)} \quad \text{(Assuming the denominator is not 0)}$$

$$\lim_{x \to 0} \frac{\sin x}{x} = 1 \qquad \qquad \qquad \lim_{x \to 0} \frac{1 - \cos x}{x} = 0$$

$$\lim_{x \to c} f(g(x)) = f(g(c)) \quad \text{(Assuming f is continuous at } \lim_{x \to c} g(x).)$$

A function f has a vertical asymptote at x = c if $\lim_{x \to c} |f(x)| = \infty$. A function g has a horizontal asymptote at y = c if $\lim_{x \to +\infty} g(x) = c$ or $\lim_{x \to -\infty} g(x) = c$.

Derivatives:

If f is differentiable at x = c, then f is continuous at x = c.

$$(x^{r})' = rx^{r-1}, \qquad (c_{1}f(x) + c_{2}g(x))' = c_{1}f'(x) + c_{2}g'(x)$$

$$(f(x)g(x))' = f'(x)g(x) + g'(x)f(x) \qquad \left(\frac{f(x)}{g(x)}\right)' = \frac{f'(x)g(x) - g'(x)f(x)}{g^{2}(x)}$$

$$(\sin x)' = \cos x \qquad (\cos x)' = -\sin x$$

$$(\tan x)' = \sec^{2} x \qquad (\cot x)' = -\csc^{2} x$$

$$(f(g(x)))' = f'(g(x))g'(x)$$

Critical points of functions over a closed interval are comprised of the interval endpoints, stationary points, and singular points.

Newton's Method for numerically solving f(x) = 0 corresponds to the iteration $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$.

Sums and integrals:

$$\sum_{i=1}^{n} i = \frac{n(n+1)}{2} \qquad \qquad \sum_{i=1}^{n} i^2 = \frac{n(n+1)(2n+1)}{6} \qquad \qquad \sum_{i=1}^{n} i^3 = \frac{(n(n+1))^2}{4}$$

If F is an antiderivative of f, then the substitution rule for integrals is,

$$\int f(g(x))g'(x)\,\mathrm{d}x = F(g(x)) + C$$

Areas and volumes:

Areas and volumes:		
Area of θ -sector of a circle	:	$\frac{\theta}{2} \times r^2 = \frac{\theta}{2} \times (\text{radius})^2$
Circular arc length sweeping out θ radians	:	$\bar{r} imes heta$
Volume of a sphere	:	$\frac{4}{3}\pi r^3 = \frac{4\pi}{3} (\text{radius})^3$
Volume of a thin cylinder	:	$A \times \Delta h = (\text{Area of base}) \times (\text{height})$
Volume of a thin shell	:	$2\pi r \times h \times \Delta r = $ (Shell circumference) × (height) × (thickness)
Crapha		

Graphs:

- Point-slope form for a line: $y y_0 = m(x x_0)$
- Equation of a circle: $(x x_0)^2 + (y y_0)^2 = r^2$