



Westfälische
Wilhelms-Universität
Münster

Contribution of EEG and MEG beamforming to the localization of interictal epileptiform dis- charges

Dissertation

Contents

1 Introduction

- 1.1 Introduction to EEG and MEG
- 1.2 Brain signals and how to measure them
- 1.3 EEG and MEG recordings
- 1.4 How head properties influence the measurements

2 Medical background

- 2.1 What is epilepsy
- 2.2 The role of EEG and MEG

3 Mathematical Background

- 3.1 The forward problem
- 3.2 The inverse problem
- 3.3 Combining EEG and MEG

4 The Beamformer approach for source reconstruction

- 4.1 A brief History
- 4.2 Formulation of the beamformer as individual spatial filter
- 4.3 Filter designs for EEG and MEG
- 4.4 The covariance estimation problem

5 Influence of the head model to the localization

- 5.1 Materials and Methods
- 5.2 Results
- 5.3 Discussion

6 A comparison of beamformer and dipole scans for somato-sensory evoked potentials and fields

- 6.1 Introduction
- 6.2 Materials and Methods
- 6.3 Results
- 6.4 Discussion

7 Beamformer for the localization of epileptic spikes

- 7.1 Materials and Methods
- 7.2 Results
- 7.3 Discussion

Contents

8 Distributional methods for spike localization

- 8.1 Introduction of simple distributional methods
- 8.2 RAMUS as a novel approach
- 8.3 Comparison between RAMUS and beamforming
 - 8.3.1 Materials and Methods
 - 8.3.2 Results
 - 8.3.3 Discussion

9 Interictal Spike Localization via Gaussian Mixture Models

- 9.1 Using Gaussian Mixture Models for clustering
- 9.2 Application for epilepsy data
 - 9.2.1 Materials and Methods
 - 9.2.2 Results
 - 9.2.3 Discussion

10 Discussion and outlook

11 Summary